



Decolourisation of Textile Dyes by Fungal Isolates

Patil Vrushali Suresh¹, Aruna Sharma², Sachin Mane³
ME Student¹, Associate Professor², Professor³
Department of Civil Engineering
D.Y.Patil College of Engineering, Akurdi, Pune, India

Abstract:

Microbial isolates of various species are studied and have found to have capacity to decolourise the textile dyes. In the present study two different fungal isolates belonging to *Trichoderma* were isolated from textile industrial effluent and were tested for their efficiency of decolourisation of textile dyes of two different colours which are commonly used in the textile industry. The efficiency of each fungus was tested for different parameters like pH, concentration of dye, and temperature. The results showed that both the species were capable of decolorizing the dyes however *Trichoderma herzianum* Rifai showed more potential of decolourisation at the pH of 5 and at 0.1 g/lit concentration of dye.

Keywords: fungi, dyes, enzymes, decolourisation etc.

I. INTRODUCTION

Textile dyes are widely used in the textile as well as in other industries as well like cosmetics, leather, printing, food, etc. these are mostly synthetic dyes. The effluent from the textile industry contains dyes as the major constituent. Most of the dyes are synthetic in nature and hence are difficult to breakdown. (Stolz 2000) Moreover these dyes are also strong to resist action of sunlight and water. These results in large quantities of dyes in the industrial waste water. The removal of colour these dyes even in small quantities are visible. These dyes reduce the penetration of sunlight and photosynthesis thus adversely affecting the aquatic flora and fauna. (Ali et al. 2010). In addition some dyes are also toxic or mutagenic and carcinogenic. (Gong et al., 2005; Nigam et al., 2000; Birhanli and Ozen, 2005; Degon et al., 2005)

II. EXPERIMENTAL METHODS

A. Materials

The dyes used in this study were BROWN 3BS, YELLOW GG and YELLOW BROWN found in abundance in the effluent water. Stock solutions of these dyes were prepared of different concentrations and the cultures were added to these solutions. Decolourisation was measured at peak absorbance wavelength of each dye using UV-VIS Spectrometer.

B. Preparation of Stock solution

Stock solutions of the three dyes were prepared by dissolving accurately measured dyes in distilled water at concentration of 500 mg/ lit. Different concentrations were prepared from this solution. (0.02 g/lit- 0.1 g/lit)

C. Fungal cultures:

Two fungal isolates belonging to *Trichoderma* viz. *Trichoderma aureoviride* Rifai and *Trichoderma herzianum* Rifai were used for the study. These fungi were isolated from the industrial effluent based on the earlier studies and literature survey.

III. EXPERIMENTAL PROCEDURE:

The effluent waste of a textile industry in Surat, Gujrat was obtained and tested for its physical parameters. 1ml of the effluent was transferred into 9 ml of distilled water in sterile test tubes. Serial dilution was done up to 10^{-7} by thorough mixing. 0.1 ml of sample from each dilution was spread on saboraraud dextrose agar plates. The petri dishes were incubated at room temperature (28°C) for 5 days. (Ponraj.M et al.) The predominant isolates were isolated and grown separately. These isolates were identified at the Agarkar research institute, Pune, India. These fungal isolates were further used for the decolourisation experiments. These were then transferred to the stock solutions in sterile conditions to avoid any kind of contamination. About 5 mm disc of the cultures were inoculated in the dye solutions and incubated for three days at 25°C.

D. Dye Decolourisation:

The experiments were performed in 250 ml flasks containing stock solutions of different concentrations (0.02-0.1 g/lit). After inoculation of fungi the flasks were incubated and were observed for decolourisation after 3, 6, and 9 days. After each period of time about 20ml sample was used and centrifuged at 7500 rpm for 20 min. then the absorbance for each concentration was measured at the wavelength of peak absorbance. The stock solutions without culture were used as control. The absorbance was taken after 3, 6 and 9 days and the percentage decolourisation was calculated. % decolourisation was determined by the formula-

$$\% \text{ Decolourization} = 100 * (C_0 - C_t) / C_t$$

Where C_0 is the initial concentration of the dye (control) and C_t is the concentration at time t.

IV. RESULTS AND DISSCUSSIONS.

The fungi are able to decolorize the dyes because of their ability of producing lignin modifying enzymes. Mostly the lacase enzymes are capable of decolourisation. The fungi used in this study were capable lacase activity which was studied in various literatures. (N.F.Ali et al.) Both the species of fungi were capable of decolorizing the dyes used on a similar

amount however the *Trichoderma aureoviride* Rifai was found to be slightly more capable of decoloring both the dyes

E. Effect of initial time of incubation on decolourisation % of dyes:

The results obtained indicated that the decolourisation efficiency of the dyes used for 3, 6, and 9 days intervals using 5 mm approx. disc of fungi culture shown in the tables 1,2, and 3 respectively, both the fungal isolates were capable of removing approximately the highest % of dyes after 9 days of incubation. The nature of the substituent on the aromatic ring has been shown to influence enzymatic oxidation. Electron donating substituents as methyl and methoxy and amino groups enhance enzymatic degradation of azo phenols while electron withdrawing substituents as chloro, nitro and hydroxyl inhibited oxidation (Zheng et al., 1999). Hydroxyl and amino groups enhance decolorization. The presence of Lip in addition to Laccase in the *Trichoderma* had positive effect on the degree of decolorization for the tested dyes. Figures 9 and 10 shows the photographs of before and after 9 days of incubation of both the dyes used.

F. Effect of initial dye concentration on % decolourisation of dyes:

The effect of initial dye concentration of dye in solution on removal of dyes was studied. We used different concentration of dyes (0.02, 0.04, 0.08 and 0.1 g/l). The removal of dyes was clearly dependent on the initial dye concentration of the solution. The highest concentration of 0.1 g/lit showed slightly greater decolorisation than the other concentrations hence as a result the same concentration was further used for evaluating the effect of pH on the % decolorisation.

G. Effect of the pH on decolourization % dyes:

Figures 5,6,7,8 show the effect of pH of the dye solution on the decolorization % of microorganisms within the range of (3–7). The results showed that the decolorization reached maximum at pH 7 for Brown color while it reached maximum at pH 5 for yellow dyes for all isolates used. Many authors stated that pH is very important for fungal growth.

Table.1. Decolourisation potential of *Trichoderma aureoviride* Rifai against different concentrations of dyes for BROWN3 BS

Concentration	3 days	6 days	9 days
0.02 g/L	3.14	6.28	26.7
0.04 g/L	18.91	25.94	27.56
0.06 g/L	21.46	27.45	31.44
0.08 g/L	26.37	28.43	35.71
0.1 g/L	23.44	33.43	56.16




Figure.1. Decolourisation potential of *Trichoderma aureoviride* Rifai against different concentrations of dyes for BROWN3 BS

Table.2. Decolourisation potential of *Trichoderma herzianum* Rifai Against different concentrations of dyes For BROWN 3BS

Concentration	3 days	6 days	9 days
0.02 g/L	12.56	26.70	36.64
0.04 g/L	19.18	25.94	30.81
0.06 g/L	23.29	28.28	31.44
0.08 g/L	25.96	28.02	31.45
0.1 g/L	26.09	30.37	46.48




Figure.2. Decolourisation potential of *Trichoderma herzianum* Rifai Against different concentrations of dyes for BROWN3 BS

Table.3. Decolourisation potential of *Trichoderma aureoviride* Rifai against different concentrations of dyes for YELLOW GG

Concentration	3 days	6 days	9 days
0.02 g/L	15.07	18.89	26.51
0.04 g/L	9.27	12.79	19.68
0.06 g/L	16.71	9.87	23.46
0.08 g/L	17	27.75	33
0.1 g/L	21.5	23.5	27




Figure.3. Decolourisation potential of *Trichoderma aureoviride* Rifai Against different concentrations of dyes for YELLOW GG

Table.4. Decolourisation potential of *Trichoderma herzianum* Rifai Against different concentrations of dyes for YELLOW GG

Concentration	3 days	6 days	9 days
0.02 g/L	9.87	15.42	25.47
0.04 g/L	5.16	11.84	22.64
0.06 g/L	10.07	12.91	20.67
0.08 g/L	8.75	13	26
0.1 g/L	5.5	16.5	23.5




Figure.4. Decolourisation potential of *Trichoderma herzianum* Rifai Against different concentrations of dyes for YELLOW GG.

Table.5. Decolourisation potential of *Trichoderma aureoviride* Rifai Against different pH values of dyes for BROWN 3BS

Ph	3 days	6 days	9 days
3	18.31	27.90	40.84
4	19.70	27.94	36.02
5	9.51	21.13	35.93
6	10.74	18.96	23.26
7	20.96	35.83	44.37




Figure.5. Decolourisation potential of *Trichoderma aureoviride* Rifai Against different pH values of dyes for BROWN 3BS

Table.6. Decolourisation potential of *Trichoderma herzianum* Rifai Against different pH values of dyes for BROWN 3BS

pH	3 days	6 days	9 days
3	54.22	69.28	82.24
4	44.83	47.79	65.94
5	51.15	57.47	60.91
6	53.12	56.21	59.30
7	43.92	53.76	55.53




Figure.6. Decolourisation potential of *Trichoderma herzianum* Rifai Against different pH values of dyes for BROWN 3BS

Table.7. Decolourisation potential of *Trichoderma aureoviride* Rifai Against different pH values of dyes for YELLOW GG.

pH	3 days	6 days	9 days
3	3.25	16.67	30.27
4	5.92	17.85	33.7
5	3.575	22.7	44.22
6	5.28	12.69	19.66
7	7.05	14.65	24.4




Figure.7. Decolourisation potential of *Trichoderma aureoviride* Rifai Against different pH values of dyes for YELLOW GG.

Table.8. Decolourisation potential of *Trichoderma herzianum* Rifai Against different pH values of dyes for YELLOW GG.

pH	3 days	6 days	9 days
3	18.31	27.90	40.84
4	19.70	27.94	36.02
5	9.51	21.13	35.93
6	10.74	18.96	23.26
7	20.96	35.83	44.37




Figure.8. Decolourisation potential of *Trichoderma herzianum* Rifai Against different pH values of dyes for YELLOW GG.

C.Effect of temperature on % decolourisation of textile dyes:

To determine the effect of temperature on the change in % decolourisation of textile dyes. The temperature was varied over a range of 10-50⁰C and the change in % decolourisation was recorded. The results showed that the maximum decolourisation was obtained in the range of 30-40⁰C maximum at 35⁰C.

Table.9. Decolourisation potential of *trichoderma aureoviride rifai* against different temperature values of dyes for YELLOW GG.

Temperature	3 days	6 days	9 days
10	6.65	12.56	25.84
20	8.25	16.85	28.02
30	12.25	21.58	35.4
35	15.41	30.22	45.21
40	8.63	24.42	35.24
50	6.25	18.96	24.56




Figure.9. Decolourisation potential of *trichoderma aureoviride rifai* against different temperature values of dyes for YELLOW GG.

Table.10. Decolourisation potential of *trichoderma herzianun rifai* against different temperature values of dyes for YELLOW GG.

Temperature	3 days	6 days	9 days
10	7.25	13.56	24.84
20	8.25	15.75	27.02
30	13.25	25.58	35.4
35	15.81	29.22	45.21
40	8.45	25.42	35.24
50	6.23	17.86	25.56




Figure.10. Decolourisation potential of *trichoderma herzianun rifai* against different temperature values of dyes for YELLOW GG.

Table.11. Decolourisation potential of *trichoderma aureoviride rifai* against different temperature values of dyes for BROWN 3BS.

Temperature	3 days	6 days	9 days
10	4.12	15.45	26.84
20	6.01	15.01	28.02
30	10.22	24.58	37.4
35	15.25	39.22	46.21
40	9.45	23.42	32.24
50	6.45	16.82	25.36




Figure.11. Decolourisation potential of trichoderma *aureoviride rifai* against different temperature values of dyes for BROWN 3BS.

Table.12. Decolourisation potential of trichoderma *herzianum rifai* against different temperature values of dyes for BROWN 3BS.

Temperature	3 days	6 days	9 days
10	5.12	15.45	23.84
20	6.01	17.01	29.02
30	11.22	22.58	38.41
35	15.45	39.72	45.23
40	10.66	25.42	32.24
50	7.22	13.82	29.10




Figure.12. Decolourisation potential of trichoderma *herzianum rifai* against different temperature values of dyes for BROWN 3BS.

D .Decolourisation of textile effluent:

The decolourisation potential of both the fungal isolates was also tested on the actual textile effluent sample.

Table.13. Decolourisation potential of Trichoderma *aureoviride Rifai* and Trichoderma *herzianum Rifai*

%Decolourisation	3 days	6 days	9 days
T.A	13.25	39.63	52.75
T.H	15.63	41.12	58.36




Figure.13. Decolourisation potential of Trichoderma *aureoviridea Rifai* and Trichoderma *herzianum Rifai*



Figure.14. Trichoderma *herzainum Rifai* and Trichoderma *aureoviridea Rifai*



Figure.14. photographs of before and after 9 days of incubation of brown 3BS dyes



Figure.15. photographs of before and after 9 days of incubation of YELLOW GG dyes.



Figure.16. photographs of before and after 9 days of incubation of Actual waste water.

V. CONCLUSION:

Fungi decolourization of dyes can be achieved by treating with Trichoderma sp. It can be stated that both fungal isolates were capable of removing the highest percent of dye colour after 9

days incubation. The isolate of *Trichoderma herzianum* Rifai produced the highest biomass accumulation after 9 days and it is the most efficient one in the removal of the reactive and acid dyes at low concentrations. The results showed that the decolorization reached maximum at pH 7 for Brown dye while it reached a maximum at pH 5 for Yellow dye for all isolates used. The results indicate that these fungal isolates can be used in conventional treatment of textile effluent to remove the dyes.

VI. REFERENCES.

- [1]. Ambrósio ST, Campos-Takaki GM.. Decolorization of reactive azo dyes by *Cunninghamella elegans* UCP 542 under co-metabolic conditions. *Bioresour Technol.* 2004;91:69–75.
- [2]. Annika A. Durvel, Arvind R. Gupta² and Sayali R. Naphade³, Decolourisation of Textile Dyes and Biological stains by Bacterial strains isolated from Industrial effluents, *Adv. Appl. Sci. Res.*, 2012, 3(5):2660-2671
- [3]. Banat IM, Nigam P, Singh D, Marchant R. Microbial decolorization of textile-dye-containing effluents: a review. *Bioresour Technol.* 1996; 58:217–27.
- [4]. Cheunbarn T, Cheunbarn S, Khumjai T. Prospects of bacterial granule for treatment of real textile industrial wastewater. *Int J Agric Biol.* 2008; 10:689–92.
- [5]. Correia VM, Stephenson T, Judd SJ. Characterization of textile wastewaters - A review. *Environ Technol.*1994; 15:917–9.
- [6]. Dubey SK, Pandey A, Bajaj AK, Misra K. Some commercial azo dyes as inhibitors of mushroom tyrosinase DOPA oxidase activity. *J Pharmacol Toxicol.* 2007;2:718–24.
- [7]. Harshad Lade, Avinash Kadam, Diby Paul, Sanjay Govindwar Biodegradation And Detoxification Of Textile Azo Dyes By Bacterial Consortium Under Sequential Microaerophilic /Aerobic Processes *EXCLI Journal* 2015;14:158-174 – ISSN 1611-2156
- [8]. J. K. Singh¹, R. Ranjan and Pranay P Pankaj, Isolation and Screening of Water Microbes for Decolourisation of Textile Dye Waste. *Current World Environment* Vol. 11(1), 296-300 (2016)
- [9]. K. Kalaiarasi, a. Lavanya, s. Amsamani and g. Bagyalakshmi, 2012, *Brazilian archives of Biology and technology*, vol.55, and 3: pp.471-476, ISSN 1516-8913 decolorization of textile dye effluent by non-viable Biomass of *Aspergillus fumigates*
- [10]. Kadam AA, Telke AA, Jagtap SS, Govindwar SP. Decolorization of adsorbed textile dyes by developed consortium of *Pseudomonas* sp. SUK1 and *Aspergillus ochraceus* NCIM-1146 under solid state fermentation. *J Hazard Mater.* 2011; 189:486–94.
- [11]. Khaled A, El Nemr A, El-Sikaily A, Abdelwahab O. Removal of Direct N Blue-106 from artificial textile dye effluent using activated carbon from orange peel: Adsorption isotherm and kinetic studies. *J Hazard Mater.* 2009; 165:100–10.
- [12]. Lade HS, Waghmode TR, Kadam AA, Govindwar SP. Enhanced biodegradation and detoxification of disperse azo dye Rubine GFL and textile industry effluent by defined fungal-bacterial consortium. *Int Biodeterior Biodegr.* 2012; 72:94–107.
- [13]. Maria Estela Silva-Stenico, Felipe D. P. Vieira, Diego B. Genuário, Caroline S. P. Silva, Luiz Alberto B. Moraes and Marli Fátima Fiore, Decolorization of Textile Dyes by Cyanobacteria, 2012, *J. Braz. Chem. Soc.*, Vol. 23, No. 10, 1863-1870.
- [14]. Meyer U. Biodegradation of synthetic organic colorants. In: Leisinger T, Cook AM, Hutter R, Nuesch J (eds): *Microbial degradation of xenobiotic and recalcitrant compounds. FEMS Symposium 12* (3rd ed.) (pp 371-85). London: Academic Press, 1981.
- [15]. N.F. Ali a, R.S.R. El-Mohamedy, Microbial decolorization of textile waste water *Journal of Saudi Chemical Society* (2012) 16, 117–123
- [16]. N. Sriram and D. Reetha, Isolation and characterization of dye degrading bacteria from textile dye Effluents, *Cent. Euro. J. Exp. Bio.*, 4 (2):5-10
- [17]. Pandey A, Singh P, Iyengar L. Bacterial decolorization and degradation of azo dyes. *Int Biodeterior Biodegr.* 2007;59:73–84.
- [18]. T. Marimuthu, S. Rrajendran, M. Manivannan, decolorisation of reactive textile dye effluent by using bacterial strain, *International Journal of Scientific and Engineering Research*, Volume 4, Issue 5, May 2013, ISSN 2229-5518.
- [19]. Wong Y, Yu J. Laccase-catalyzed decolorization of synthetic dyes. *Water Res.* 1999;33:3512–20.